


Scattering amplitudes and hidden symmetries in supersymmetric gauge theory

Jan Plefka

Institut für Physik, Humboldt-Universität zu Berlin
und Institut für Theoretische Physik, ETH Zürich



ETH zürich




Scattering amplitudes and hidden symmetries in maximally supersymmetric gauge theory

Plan:

- 1 Introduction: Quantum field theory
- 2 Symmetries
- 3 Supersymmetric gauge field theory
- 4 Scattering amplitudes in gauge theories and on-shell methods
- 5 Symmetries of scattering amplitudes
- 6 Generalized unitarity
- 7 (String-gauge theory duality)

Mathematical framework of particle physics: QFT

- **Quantum Field Theory**: **Relativistic** many particle quantum theory
- Describes scattering processes in accelerators



Perturbative description: Series expansion in $g \ll 1$ g : Coupling constant

- **Feynman diagrams**: Describe particle propagation & interactions
- **Symmetries** play central role:
 - Determine possible particles & their interactions
 - Can severely constrain results for observables
- Exact analytic methods beyond perturbation theory are sparse
- Desirable to advance our fundamental understanding of quantum field theory

The Standard Model of Particle Physics

Three fundamental forces described by

Gauge Field Theories

[1955,1971]

Forces:

SU(3) × SU(2) × U(1)

≐ Gauge Field Theories

Electromagnetism (photons)

Weak Force (W & Z bosons)

Strong Force (gluons) ≐ Quantum Chromodynamics (QCD)

SU(N) Gauge Field Theory: Fields are $N \times N$ matrices: $\mathbf{A}_\mu^{\text{SU}(2)}(x) = \begin{pmatrix} Z & W^+ \\ W^- & -Z \end{pmatrix}$

Leptons	Quarks	Vector bosons	Scalar
e, ν_e	u, d	A_μ	
μ, ν_μ	s, c	W^\pm, Z	Higgs
τ, ν_τ	t, b	A_μ^a	

Gravity is not contained!

Spectrum:

Symmetries

Symmetries

Symmetries lie at the heart of our understanding of physics. They constrain or even determine physical theories and their observables.

- Mathematically symmetry transformations form a group

$$G_1 \circ G_2 = G_3 \quad \{G_i, \mathbb{1}, G_i^{-1}\} \in \text{group}$$

- Continuous transf.: Lie group $G(\phi) = e^{i\phi^a \hat{J}_a}$ \hat{J}_a : Generator $\phi^a \in \mathbb{R}$
- Group property entails commutation relations

$$[\hat{J}_a, \hat{J}_b] = i f_{ab}^c \hat{J}_c \quad \text{Lie algebra} \quad a, b, c = 1, \dots, \dim(g)$$

- Symmetries can be obvious or hidden
- Example:** Rotations and translations

$$R(\vec{\phi}) = e^{i\vec{\phi} \cdot \hat{L}}$$

\hat{L} : Angular momentum

$$T(\vec{a}) = e^{i\vec{a} \cdot \hat{P}}$$

\hat{P} : Momentum

$$[L_i, L_j] = i\hbar \epsilon_{ijk} L_k \quad [P_i, P_j] = 0 \quad [L_i, P_k] = i\hbar \epsilon_{ijk} P_k$$

Symmetries

Symmetries lie at the heart of our understanding of physics. They constrain or even determine physical theories and their observables.

- Mathematically symmetry transformations form a group

$$G_1 \circ G_2 = G_3 \quad \{G_i, \mathbb{1}, G_i^{-1}\} \in \text{group}$$

- Continuous transf.: Lie group $G(\phi) = e^{i\phi^a \hat{J}_a}$ \hat{J}_a : Generator $\phi^a \in \mathbb{R}$
- Group property entails commutation relations

$$[\hat{J}_a, \hat{J}_b] = if_{ab}^c \hat{J}_c \quad \text{Lie algebra} \quad a, b, c = 1, \dots, \dim(g)$$

- Symmetries can be obvious or hidden
- Example:** Rotations and translations


$$R(\vec{\phi}) = e^{i\vec{\phi} \cdot \hat{\vec{L}}} \qquad T(\vec{a}) = e^{i\vec{a} \cdot \hat{\vec{P}}}$$

\vec{L} : Angular momentum \vec{P} : Momentum

$$[L_i, L_j] = i\hbar \epsilon_{ijk} L_k \qquad [P_i, P_j] = 0 \qquad [L_i, P_k] = i\hbar \epsilon_{ijk} P_k$$

Hidden symmetries: The Hydrogen atom

- Hamiltonian
$$H = \frac{\vec{p}^2}{2m} - \frac{k}{r}$$
- Rotational symmetry: $[H, L_i] = 0 \Rightarrow H |n, l, m\rangle = E_{n,l} |n, l, m\rangle$
- Hidden symmetry in H-atom: **Pauli-Lenz vector**

$$\vec{A} = \frac{1}{2}(\vec{p} \times \vec{L} - \vec{L} \times \vec{p}) - m k \frac{\vec{r}}{r}$$


- Conserved quantity: $[H, A_i] = 0$
- Algebra:

$$[A_i, A_j] = -i \frac{2\hbar}{m} H L_k, \quad [L_i, A_j] = i\hbar \epsilon_{ijk} A_k, \quad [L_i, L_j] = i\hbar \epsilon_{ijk} L_k$$

- Closes on eigenspace \mathcal{H}_E of fixed energy eigenvalue E .
- Operator algebra determines spectrum ($\hat{=}$ representation theory of $SU(2)$)

$$E_n = -\frac{mk^2}{2\hbar^2} \frac{1}{n^2} \quad (\text{degeneracy } n^2)$$

Fundamental symmetry of QFT: Poincaré group

Poincare symmetry: Lorentz transformations + translations in space & time

$$M_{\mu\nu}, P_\mu$$

\vec{L}
rotations

$$\vec{K}$$

boosts

$$P_\mu$$

$$L_i = \frac{1}{2} \epsilon_{ijk} M_{jk}$$

$$K_i = M_{0i}$$

$$\mu, \nu, \dots = \{0, i\}$$

• Representations

spin	field	example
0	scalar $\phi(x)$	Higgs
1/2	left handed spinor $\chi_\alpha(x)$	leptons, quarks
1/2	right handed spinor $\bar{\psi}_{\dot{\alpha}}(x)$	leptons, quarks
1	vector $A_\mu(x)$	photon, gauge bosons
3/2	$\psi_\mu^\alpha(x)$	gravitino (Rarita-Schwinger field)
2	$h_{\mu\nu}(x)$	graviton

- Massless fields only have helicity $h = \frac{\vec{p} \cdot \vec{S}}{|\vec{p}|}$ states $h = \pm s$

Extension I: Conformal symmetry

- Physical theories without intrinsic mass scale ($\hat{=}$ massless theories or at very high energies) have an enlarged space-time symmetry: **Conformal symmetry**
- Angle preserving transformations: Dilatations and inversions

Dilatation transf.: $d: x^\mu \rightarrow \kappa x^\mu \quad \kappa \in \mathbb{R}$

Special conformal transf.: $k^\mu = I \circ p^\mu \circ I$ with I : Inversion $x^\mu \rightarrow \frac{x^\mu}{x^2}$

- Conformal group is $\mathfrak{so}(2, 4)$ with algebra:

$$[K_\mu, P_\nu] = 2i(\eta_{\mu\nu}D - M_{\mu\nu}), \quad [D, P_\mu] = iP_\mu, \quad [D, K_\mu] = -iK_\mu, \\ [K_\rho, M_{\mu\nu}] = i(\eta_{\rho\mu}K_\nu - \eta_{\rho\nu}K_\mu) \quad \& \text{ Poincaré algebra}$$

- Examples:

Maxwell's theory	$\mathcal{L} = \frac{1}{4}F_{\mu\nu}F^{\mu\nu}, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$
$\lambda\phi^4$ theory	$\mathcal{L} = \frac{1}{2}(\partial_\mu\phi)^2 - \lambda\phi^4$
Standard model up to Higgs mass term	$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\bar{\psi}\not{D}\psi + \psi_i Y_{ij}\psi_j \phi \\ + D_\mu\phi ^2 - \lambda \phi ^4 - m^2 \phi ^2$

Extension I: Conformal symmetry

- Physical theories without intrinsic mass scale ($\hat{=}$ massless theories or at very high energies) have an enlarged space-time symmetry: **Conformal symmetry**
- Angle preserving transformations: Dilatations and inversions

Dilatation transf.: $d: x^\mu \rightarrow \kappa x^\mu \quad \kappa \in \mathbb{R}$

Special conformal transf.: $k^\mu = I \circ p^\mu \circ I$ with I : Inversion $x^\mu \rightarrow \frac{x^\mu}{x^2}$

- Conformal group is $\mathfrak{so}(2, 4)$ with algebra:

$$[K_\mu, P_\nu] = 2i(\eta_{\mu\nu}D - M_{\mu\nu}), \quad [D, P_\mu] = iP_\mu, \quad [D, K_\mu] = -iK_\mu, \\ [K_\rho, M_{\mu\nu}] = i(\eta_{\rho\mu}K_\nu - \eta_{\rho\nu}K_\mu) \quad \& \text{ Poincaré algebra}$$

- Examples:

Maxwell's theory	$\mathcal{L} = \frac{1}{4}F_{\mu\nu}F^{\mu\nu}, \quad F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$
$\lambda\phi^4$ theory	$\mathcal{L} = \frac{1}{2}(\partial_\mu\phi)^2 - \lambda\phi^4$
Standard model up to Higgs mass term	$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\bar{\psi}\not{D}\psi + \psi_i Y_{ij}\psi_j \phi \\ + D_\mu\phi ^2 - \lambda \phi ^4 - m^2 \phi ^2$

Extension II: Supersymmetry

Supersymmetry is a unique extension of space-time symmetries [1971,1974]

“Square root” of the momentum:

$$\{Q_\alpha, \bar{Q}_{\dot{\alpha}}\} = 2(\sigma^\mu)_{\alpha\dot{\alpha}} P_\mu$$

- Graded Lie algebra: Generators Q_α & $\bar{Q}_{\dot{\alpha}}$ are fermionic.
Obey Super-Poincaré algebra with generators $\{M_\mu, P_\mu; Q_\alpha, \bar{Q}_{\dot{\alpha}}\}$
- Relates **bosons** and **fermions**:

$$\bar{Q}_{\dot{\alpha}} |\text{spin} = s\rangle = |\text{spin} = s + 1/2\rangle$$

- SUSY:

Boson	\longleftrightarrow	Fermion
Gluon	\longleftrightarrow	Gluino

Superpartners are degenerate in all quantum numbers (mass, charge, ...)

- Extended supersymmetry**: Can have more than one set of supercharges

$\rightarrow Q_\alpha^A$ & $\bar{Q}_{\dot{\alpha}A}$ with $A = 1, \dots, \mathcal{N}$:

B_1	\longleftrightarrow	F_1
B_2	\longleftrightarrow	F_2

Gluon \longleftrightarrow \mathcal{N} **Gluinos**

- Maximal SUSY**: $\mathcal{N} = 4$ spin-range $\{-1, -1/2, 0, 1/2, 1\}$

Gauge field theory

Gauge Field Theory (or Yang-Mills-Theory)

- Builds upon **internal** (non-space-time) symmetry
- **SU(N) Gauge theory:** [1954]

Generalization of Maxwell's theory of electromagnetism: $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$

Vector potential now $N \times N$ hermitian matrix: $(\mathbf{A}_\mu)_{ab}(x) \quad a,b=1,\dots,N$

- **Local** gauge symmetry: $\mathbf{A}_\mu(x) \rightarrow \mathbf{U} \mathbf{A}_\mu \mathbf{U}^\dagger + \frac{i}{g} \mathbf{U} \partial_\mu \mathbf{U}^\dagger \quad \partial_\mu = \frac{\partial}{\partial x^\mu}$
with $\mathbf{U} \in SU(N)$, i.e. unitary $N \times N$ matrix, $\mathbf{U} \mathbf{U}^\dagger = 1$

- Invariant action

$$S_{\text{YM}} = \frac{1}{4} \int d^4x \text{Tr}(\mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu}) \quad \mathbf{F}_{\mu\nu} = \partial_\mu \mathbf{A}_\nu - \partial_\nu \mathbf{A}_\mu + i g [\mathbf{A}_\mu, \mathbf{A}_\nu]$$

g: Coupling constant.



- $N = 1$: Maxwell theory!

$\mathcal{N} = 4$ super Yang-Mills theory

Can we have everything?


- Poincaré symmetry \rightarrow relativistic QFT
- Conformal symmetry \rightarrow scale-invariant
- Maximal supersymmetry ($\mathcal{N} = 4$)
- $SU(N)$ local gauge symmetry (with $N \rightarrow \infty$)

\Rightarrow $\mathcal{N} = 4$ SYM

A_μ	1 Gluon	spin=1		(= same as in QCD)
ψ_α^A	4 Gluinos	spin=1/2		(= cousin of the quarks)
ϕ_I	6 Scalars	spin=0	

$$\mathcal{L}_{\text{SYM}} = \frac{N}{\lambda} \text{Tr} \left[\frac{1}{4} F_{\mu\nu}^2 + \frac{1}{2} (D_\mu \Phi_I)^2 - \frac{1}{4} [\Phi_I, \Phi_J][\Phi_I, \Phi_J] + \bar{\psi} \not{D} \psi + \psi^A \psi^B \Phi_{AB} + h.c. \right]$$

Interactions:



All fields $N \times N$ matrices. In $N \rightarrow \infty$ (planar) limit: One parameter $\lambda = g^2 N$

The simplest quantum field theory

$\mathcal{N} = 4$ SYM has remarkably rich properties:

- Uniquely determined by g_{YM} & N , exactly scale invariant at any coupling, no UV divergences $\Rightarrow g_{YM} = \text{const}$ [1980's]
- Dual to string theory \rightarrow AdS/CFT correspondence. [1997]
Strong coupling limit ($\lambda = g^2 N^2 \rightarrow \infty$): Classical string on $AdS_5 \times S^5$.
- Appears to be integrable in $N \rightarrow \infty$ limit: [since 2003]
 - Exact results for two-point correlation functions $\langle \mathcal{O}(x)\mathcal{O}(y) \rangle = (x - y)^{2\Delta + \gamma(\lambda)}$
 - Hidden symmetries beyond super-conformal group: Yangian algebra
 - Deep mathematical understanding of scattering amplitudes


Renders model an ideal theoretical laboratory to study gauge theories (and string theory)!

Could be the first exactly solvable interacting 4d QFT.

- \Rightarrow Non-physical! But possible starting point for novel perturbative approach.
- \Rightarrow Already now application to massless QCD exist.

Scattering amplitudes


Scattering amplitudes





$$\mathcal{A}_n(\{p_i, h_i\}) = \text{probability amplitude for scattering process}$$

Central quantum field theory prediction for collider experiments

Computed via Feynman diagrams:

Propagator  $= \frac{\delta^{ab} \eta_{\mu\nu}}{k^2 + i\epsilon}$ (gluons)

Vertices  $= g f^{abc} \left[(q-r)_\mu \eta_{\nu\rho} + (r-p)_\nu \eta_{\rho\mu} + (p-q)_\rho \eta_{\mu\nu} \right]$



$$= -ig^2 \left[f^{abe} f^{cde} (\eta_{\mu\rho} \eta_{\nu\sigma} - \eta_{\mu\sigma} \eta_{\nu\rho}) + f^{ace} f^{bde} (\eta_{\mu\sigma} \eta_{\rho\nu} - \eta_{\mu\nu} \eta_{\rho\sigma}) + f^{ade} f^{bce} (\eta_{\mu\nu} \eta_{\sigma\rho} - \eta_{\mu\rho} \eta_{\sigma\nu}) \right]$$




Feynman diagrammatics

Task:

- Draw all Feynman diagrams contributing to a given process
- Integrate over all intermediate (off-shell) momenta $\int d^{4-2\epsilon}l$ imposing momentum conservation $\delta^{(4)}(\sum_i p_i)$ at each vertex
- $\mathcal{A}_n = \sum$ all diagrams

Can rapidly get out of hand: (even at tree-level)



number of external gluons	4	5	6	7	8	9	10
number of diagrams	4	25	220	2485	34300	559405	10525900

Simplicity of the result

When expressed in right variables the result is remarkably simple:

[Parke,Taylor]

$$\mathcal{A}_5(1^\pm, 2^+, 3^+, 4^+, 5^+) = 0$$

$$\mathcal{A}_5(1^-, 2^-, 3^+, 4^+, 5^+) = \frac{\langle 12 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 45 \rangle \langle 51 \rangle}$$

$$\mathcal{A}_5(1^-, 2^+, 3^-, 4^+, 5^+) = \frac{\langle 13 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 45 \rangle \langle 51 \rangle}$$

(all others from cyclicity and parity)

Spinor helicity:

$$\boxed{p^\mu \rightarrow p^{\alpha\dot{\alpha}} = \bar{\sigma}_\mu^{\alpha\dot{\alpha}} p^\mu = \lambda^\alpha \tilde{\lambda}^{\dot{\alpha}}} \quad (\text{makes } p^\mu p_\mu = 0 \text{ manifest})$$

$$\lambda^\alpha = \frac{1}{\sqrt{p^0 + p^3}} \begin{pmatrix} p^0 + p^3 \\ p^1 + ip^2 \end{pmatrix}, \quad \tilde{\lambda}^{\dot{\alpha}} = (\lambda^\alpha)^\dagger, \quad \langle ij \rangle = \epsilon_{\alpha\beta} \lambda_i^\alpha \lambda_j^\beta$$

What is the reason for this simplicity?

- Hidden symmetries (\rightarrow hidden super-conformal invariance & more)
- Analytic structure of the amplitude (\rightarrow factorization, soft & collinear limits)

Simplicity of the result

When expressed in right variables the result is remarkably simple:

[Parke,Taylor]

$$\mathcal{A}_5(1^\pm, 2^+, 3^+, 4^+, 5^+) = 0$$

$$\mathcal{A}_5(1^-, 2^-, 3^+, 4^+, 5^+) = \frac{\langle 12 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 45 \rangle \langle 51 \rangle}$$

$$\mathcal{A}_5(1^-, 2^+, 3^-, 4^+, 5^+) = \frac{\langle 13 \rangle^4}{\langle 12 \rangle \langle 23 \rangle \langle 34 \rangle \langle 45 \rangle \langle 51 \rangle}$$

(all others from cyclicity and parity)

Spinor helicity:

$$\boxed{p^\mu \rightarrow p^{\alpha\dot{\alpha}} = \bar{\sigma}_\mu^{\alpha\dot{\alpha}} p^\mu = \lambda^\alpha \tilde{\lambda}^{\dot{\alpha}}} \quad (\text{makes } p^\mu p_\mu = 0 \text{ manifest})$$


$$\lambda^\alpha = \frac{1}{\sqrt{p^0 + p^3}} \begin{pmatrix} p^0 + p^3 \\ p^1 + ip^2 \end{pmatrix}, \quad \tilde{\lambda}^{\dot{\alpha}} = (\lambda^\alpha)^\dagger, \quad \langle ij \rangle = \epsilon_{\alpha\beta} \lambda_i^\alpha \lambda_j^\beta$$

What is the reason for this simplicity?

- Hidden symmetries (\rightarrow hidden super-conformal invariance & more)
- Analytic structure of the amplitude (\rightarrow factorization, soft & collinear limits)

Basic problem of Feynman diagrammatic approach

In Feynman graph techniques one sums and integrates over non-physical terms:

$$\int \frac{d^3 p dE}{(2\pi)^4}$$

$$E^2 - \vec{p}^2 \neq m^2$$

Internal states are off-shell, violate mass-shell condition

Similarly individual diagrams are gauge variant, but final result is gauge invariant!

On-shell approaches:

Since 2005 tremendous progress in our understanding of scattering amplitudes based on on-shell formulations:

- On-shell recursion relations ✓
- Hidden symmetries ✓
- Generalized unitarity ✓
- Twistors & the Grassmannian & Integrability ✗

The $\mathcal{N} = 4$ SYM theory has been instrumental in this progress!

Britto-Cachazo-Feng-Witten (BCFW) recursion

- Idea: Complexify momenta but stay on-shell $z \in \mathbb{C}$

$$p_1 \rightarrow \hat{p}_1(z) = \lambda_1 (\tilde{\lambda}_1 - z \tilde{\lambda}_n) \quad p_n \rightarrow \hat{p}_n(z) = (\lambda_n + z \lambda_1) \tilde{\lambda}_n$$

Obeys $\hat{p}_i(z)^2 = 0$ and $\hat{p}_1(z) + p_2 + \dots + p_{n-1} + \hat{p}_n(z) = 0$.


$$\mathcal{A}_n \rightarrow \mathcal{A}_n(z) \quad \mathcal{A}_n(z=0) = \sum \text{Res } \mathcal{A}_n(z_i)$$

- Cauchy's theorem yields recursive relation for on-shell amplitudes

$$\mathcal{A}_n = \sum_{i,h} \mathcal{A}_{i+1}^h \frac{1}{P_i^2} \mathcal{A}_{n-i+1}^{-h}$$


- “Atoms” are the 3-point amplitudes: $\mathcal{A}_3(i^-, j^-) = \frac{\langle ij \rangle}{\langle 12 \rangle \langle 23 \rangle \langle 31 \rangle}$

- Example:




$\mathcal{N} = 4$ SYM: Superamplitudes and Super-BCFW recursion

- Consider **super** momentum-space using **4 anti-commuting coordinates** η^A :

- Define superamplitudes \mathbb{A}_n in this formal space:

[Nair]



$$\mathbb{A}_n = \frac{\delta^{(4)}(\sum_i p_i) \delta^{(8)}(\sum_i q_i)}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle} \mathcal{P}_n(\{\lambda_i, \tilde{\lambda}_i, \eta_i\})$$

Superamplitudes package all gluon-gluino-scalar amplitudes.

- Super-BCFW recursion exists:

[Brandhuber, Heslop, Travaglini][Arkani-Hamed, Cachazo, Cheung, Kaplan]

$$\mathbb{A}_n(1, \dots, n) = \sum_{i=3}^{n-1} \int d^4 \eta_{\hat{P}_i} \mathbb{A}_i^L(\hat{1}, \dots, -\hat{P}_i) \frac{1}{P_i^2} \mathbb{A}_{n-i+2}^R(\hat{P}_i, \dots, \hat{n})$$

- May be solved analytically!

⇒ **All** tree-amplitudes in $\mathcal{N} = 4$ SYM known in analytic form.


[Drummond, Henn]

Application to massless QCD

Use gluon-gluino amplitudes from $\mathcal{N} = 4$ SYM to construct compact analytic formulae for all n -point tree-level gluon-quark ($g^{n-2l}(q\bar{q})^l$) amplitudes with $l \leq 4$:

[Dixon,Henn,JP,Schuster]

Needs to suppress intermediate production of scalars



Leads to numerically fast and stable results


[Badger,Biedermann,Hackl,JP,Schuster,Uwer]

⇒ Mathematica package GGT available [Dixon,Henn,JP,Schuster]

⇒ Formulae are being used for cross section computations of LHC processes today!

[BlackHat collaboration]

(N-2) gluon 2 quark amplitudes



Symmetries of scattering amplitudes

- Superconformal symmetry of $\mathcal{N} = 4$ SYM constrains superamplitudes

$$\mathbb{A}_n^{\text{tree}} = \frac{\delta^{(4)}(\sum_i p_i) \delta^{(8)}(\sum_i q_i)}{\langle 12 \rangle \langle 23 \rangle \dots \langle n1 \rangle} \mathcal{P}_n(\lambda_i, \tilde{\lambda}_i, \eta_i)$$

- Obvious symmetries:

$$p^{\alpha\dot{\alpha}} = \sum_{i=1}^n \lambda_i^\alpha \tilde{\lambda}_i^{\dot{\alpha}} \quad q^{\alpha A} = \sum_{i=1}^n \lambda_i^\alpha \eta_i^A \quad \Rightarrow \quad p^{\alpha\dot{\alpha}} \mathbb{A}^{\text{tree}} = 0 = q^{\alpha A} \mathbb{A}^{\text{tree}}$$

explains vanishing of $A_n(1^\pm, 2^+, \dots, n^+)$

- Less obvious symmetries

[Witten]

$$k_{\alpha\dot{\alpha}} = \sum_{i=1}^n \frac{\partial}{\partial \lambda_i^\alpha} \frac{\partial}{\partial \tilde{\lambda}_i^{\dot{\alpha}}} \quad s_{\alpha A} = \sum_{i=1}^n \frac{\partial}{\partial \lambda_i^\alpha} \frac{\partial}{\partial \eta_i^A} \quad \Rightarrow \quad k_{\alpha\dot{\alpha}} \mathbb{A}^{\text{tree}} = 0 = s_{\alpha A} \mathbb{A}^{\text{tree}}$$

explains form of $A_n(1^-, 2^-, 3^+ \dots, n^+)$

- We have super-conformal invariance of tree-amplitudes (32+32 generators):

$$J^a \mathbb{A}_n^{\text{tree}} = 0 \quad \text{with} \quad J^a \in \{p, k, \bar{m}, m, d, r, q, \bar{q}, s, \bar{s}, c_i\}$$

Infinite dimensional hidden symmetry

Tree superamplitudes are invariant under additional hidden symmetry (as in H-atom)

[Drummond,Henn,Korchinsky,Sokatchev][Drummond,Henn,JP]

- Mathematical structure: Yangian algebra $Y[\mathfrak{psu}(2, 2|4)]$

[Drinfeld]

$$J^a = \sum_{i=1}^n J_i^a \quad (\text{level } 0) \quad J_{(1)}^a = f^a_{bc} \sum_{i<j}^n J_i^b J_j^c \quad (\text{level } 1)$$

An ∞ -dim non-local symmetry algebra $J_{(n)}^a \quad n = 0, 1, 2, \dots$

$$\begin{aligned} [J^a, J^b] &= i f^{ab}{}_c J^c \\ [J^a, J_{(1)}^b] &= i f^{ab}{}_c J_{(1)}^c \\ [J_{(1)}^a, J_{(1)}^b] &= i f^{ab}{}_c J_{(2)}^c + g_{ab}(J^a, J_{(1)}^a) \end{aligned}$$

$$\boxed{J_{(n)}^a \mathbb{A}_n^{\text{tree}} = 0} \quad \forall n \quad [\text{Drummond,Henn,JP}]$$

- Signature of integrable field theory. Explains simplicity of $\mathbb{A}_n^{\text{tree}}$

\Leftrightarrow Determines form of $\mathbb{A}_n^{\text{tree}}$ [Bargheer,Beisert,McLoughlin,Loebbert,Galleas]

- AdS/CFT: T-duality of dual string theory. [Alday,Maldacena][Beisert,Ricci,Tseytlin][Berkovits,Maldacena]

- **Tree level** scattering amplitudes $\hat{=}$ Sum of Yangian invariants

$$|\Psi\rangle_{n,p} = \mathcal{B}_{i_1 j_1}(u_1) \dots \mathcal{B}_{i_p j_p}(u_p) |\mathbf{0}\rangle$$

Based on methods of “quantum inverse scattering method” \Rightarrow Towards an **algebraic S-Matrix** [Chicherin,Derkachov,Kirschner] [Kanning,Lukowski,Staudacher][Broedel,de Leuw,Rosso]


- **Loop level scattering amplitudes:**

- IR divergencies break conformal symmetry in a controlled way: **Conformal anomaly** [Drummond,Henn,Korchemsky,Sokatchev]
- **Deformed** Yangian symmetry a 1-loop level [Beisert,Henn,McLoughlin,JP]
- All loop **integrands** are Yangian invariant and constructible via loop level BCFW recursion [Arkani-Hamed,Bourjaily,Cachazo,Caron-Huot,Trnka]
- And more . . .

Generalized unitarity

- On-shell methods also constructive at 1-loop (NLO-order) [Bern,Dixon,Dunbar,Kosower]
- General 1-loop amplitude may be decomposed in basis integrals

[Passarino,Veltman][Ossola,Papadopoulos,Pittau][Giele,Kunszt,Melnikov]



- In $\mathcal{N} = 4$ SYM: Only box integrals occur due to hidden symmetry.

$$A_n^{1\text{-loop}} = \sum_i c_i \text{Box}_i$$


- Find c_i by putting internal propagators on-shell

[Bern,Dixon,Kosower,Smirnov]

$$c_i = \begin{array}{c} \text{Diagram of a box integral with dashed internal lines and external lines} \\ \text{with a blue arrow pointing to a vertex and the equation } E^2 - \vec{p}^2 = 0 \end{array} = \frac{1}{2} \sum_{l_{\pm}} A_1^{\text{tree}}(l_{\pm}) A_2^{\text{tree}}(l_{\pm}) A_3^{\text{tree}}(l_{\pm}) A_4^{\text{tree}}(l_{\pm})$$

State of the art

Known MHV amplitudes: $A_n(1^-, 2^-, 3^+, \dots, n^+)$ in $\mathcal{N} = 4$ SYM



Gauge - String Theory Duality

Dual string theory in a nut-shell

- **Idea:** Replace particle by extended 1d object: **string**



- Quantum mechanics of a relativistic string in **flat space-time** $\mathbb{R}^{1,d-1}$:



Graviton



Gauge boson



Matter particle


- Oscillation spectrum $\hat{=}$ spectrum of “elementary particles”
- **Strings must** propagate in $d=9+1$. Theory depends on **background geometry**
- **Now:** Take **curved space-time geometry with boundary:** Anti-de-Sitter₅ \times 5-Sphere
- Motivation for a duality to gauge fields . . .

Quantum gravity in a box

- Space-time with negative curvature: anti-de-Sitter space (AdS_d)

[Willem de Sitter, 1872-1934]


- AdS_5 is (4+1)-dimensional space-time with boundary of geometry $\mathbb{R}^{1,3}$



- String theory well defined on $AdS_5 \times M_5$, e.g. $M_5 = S^5$ (5d-sphere).
- **Quantization** of strings on $AdS_5 \times S^5$ **unsolved!**
- Isometry group of $AdS_5 \hat{=} \text{conformal group in 4d}$

The String-Gauge Theory (or AdS/CFT) duality [Maldacena, 1997]


- **Holographic duality:** Strings in $AdS_5 \times S^5$ are dual to $\mathcal{N} = 4$ SYM



- Hidden symmetry of gauge theory may be understood as intrinsic symmetry (T-duality) of dual $AdS_5 \times S^5$ superstring theory

Summary

Field combines a multitude of areas in theoretical and mathematical physics:



⇒ Intellectually rich and fascinating research area with “real physics” applications!

Thank you for your attention

Literature:

Bern, Dixon, Kosower, Scientific American 2012

Beisert et. al. „Review of AdS/CFT integrability“, Lett.Math.Phys.99

Ellis, Kunszt, Melnikov, Zanderighi, Phys. Rep. 518 (2012)

Henn & Plefka, „Scattering Amplitudes in Gauge Theories“
LNP 883, Springer

