

# The quark-antiquark potential in $N=4$ SYM from an open spin-chain

Nadav Drukker




Based on [arXiv:1105.5144](https://arxiv.org/abs/1105.5144) - N.D. and V. Forini  
[arXiv:1203.1617](https://arxiv.org/abs/1203.1617) - N.D.

See also [arXiv:1203.1913](https://arxiv.org/abs/1203.1913) - D. Correa, J. Maldacena and A. Sever

International Symposium Ahrenshoop

August 28, 2012



## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.

## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.
- Such an object exists also in  $\mathcal{N} = 4$  SYM.
  - The Wilson loop calculates the potential between two W-bosons arising from a Higgs mechanism.

## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.
- Such an object exists also in  $\mathcal{N} = 4$  SYM.
  - The Wilson loop calculates the potential between two W-bosons arising from a Higgs mechanism.
- Explicit calculations at weak and at strong coupling:

$$V(L, \lambda) = \begin{cases} -\frac{\lambda}{4\pi L} + \frac{\lambda^2}{8\pi^2 L} \ln \frac{T}{L} + \dots & \lambda \ll 1 \\ \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L} \left( 1 - \frac{1.3359 \dots}{\sqrt{\lambda}} + \dots \right) & \lambda \gg 1 \end{cases}$$

## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.
- Such an object exists also in  $\mathcal{N} = 4$  SYM.
  - The Wilson loop calculates the potential between two W-bosons arising from a Higgs mechanism.
- Explicit calculations at weak and at strong coupling:

$$V(L, \lambda) = \begin{cases} -\frac{\lambda}{4\pi L} + \frac{\lambda^2}{8\pi^2 L} \ln \frac{1}{\lambda} + \dots & \lambda \ll 1 \\ \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L} \left( 1 - \frac{1.3359 \dots}{\sqrt{\lambda}} + \dots \right) & \lambda \gg 1 \end{cases}$$

## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.
- Such an object exists also in  $\mathcal{N} = 4$  SYM.
  - The Wilson loop calculates the potential between two W-bosons arising from a Higgs mechanism.
- Explicit calculations at weak and at strong coupling:

$$V(L, \lambda) = \begin{cases} -\frac{\lambda}{4\pi L} + \frac{\lambda^2}{8\pi^2 L} \ln \frac{1}{\lambda} + \dots & \lambda \ll 1 \\ \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L} \left( 1 - \frac{1.3359 \dots}{\sqrt{\lambda}} + \dots \right) & \lambda \gg 1 \end{cases}$$

- Recently  $O(\lambda^3)$  was calculated.
- Hard to guess how to connect these two regimes.

[ Correa, Henn  
Maldacena, Sever ]

## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.
- Such an object exists also in  $\mathcal{N} = 4$  SYM.
  - The Wilson loop calculates the potential between two W-bosons arising from a Higgs mechanism.
- Explicit calculations at weak and at strong coupling:

$$V(L, \lambda) = \begin{cases} -\frac{\lambda}{4\pi L} + \frac{\lambda^2}{8\pi^2 L} \ln \frac{1}{\lambda} + \dots & \lambda \ll 1 \\ \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L} \left( 1 - \frac{1.3359\dots}{\sqrt{\lambda}} + \dots \right) & \lambda \gg 1 \end{cases}$$

- Recently  $O(\lambda^3)$  was calculated.
- Hard to guess how to connect these two regimes.
- Can we do any better?

[ Correa, Henn  
Maldacena, Sever ]

## Introduction and motivation

- One of the most fundamental quantities in a quantum field theory is the potential between charged particles.
- In gauge theories this is captured by a long rectangular Wilson loop, or a pair of antiparallel lines.
- Such an object exists also in  $\mathcal{N} = 4$  SYM.
  - The Wilson loop calculates the potential between two W-bosons arising from a Higgs mechanism.

- Explicit calculations at weak and at strong coupling:

$$V(L, \lambda) = \begin{cases} -\frac{\lambda}{4\pi L} + \frac{\lambda^2}{8\pi^2 L} \ln \frac{1}{\lambda} + \dots & \lambda \ll 1 \\ \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L} \left( 1 - \frac{1.3359 \dots}{\sqrt{\lambda}} + \dots \right) & \lambda \gg 1 \end{cases}$$

- Recently  $O(\lambda^3)$  was calculated. [ Correa, Henn  
Maldacena, Sever ]
- Hard to guess how to connect these two regimes.
- Can we do any better?
- Shouldn't **integrability** allow us to calculate this for all values of the coupling (in the **planar** approximation)?



## Outline

- Introduction and motivation
- Wilson loops
  - Cusp anomalous dimensions and the quark-antiquark potential
  - Local operator insertions
- Generalize quark-antiquark potential in  $\mathcal{N} = 4$  SYM
  - Perturbative calculation
  - String calculation
  - Expansions in small angles
- Wilson loops and integrability
  - Operator insertions and open spin-chains
  - All loop reflection matrix and a twist
  - Wrapping effects and the quark-antiquark potential

## Wilson loops

- In any gauge theory one can define Wilson loop operators

$$W = \text{Tr} \mathcal{P} \exp \left[ \oint i A_\mu \dot{x}^\mu ds \right]$$

- Can be defined for an arbitrary curve in spacetime.

# Wilson loops

- In any gauge theory one can define Wilson loop operators


$$W = \text{Tr} \mathcal{P} \exp \left[ \oint i A_\mu \dot{x}^\mu ds \right]$$

- Can be defined for an arbitrary curve in spacetime.
- For a pair of antiparallel lines

$$\langle W \rangle \approx \exp \left[ - T V(L, \lambda) \right]$$

- The potential behaves like

$$V(L, \lambda) = \begin{cases} g(\lambda) & \text{screening} \\ \frac{f(\lambda)}{L} & \text{conformal} \\ \alpha' L & \text{confining} \end{cases}$$



# Wilson loops

- In any gauge theory one can define Wilson loop operators

$$W = \text{Tr} \mathcal{P} \exp \left[ \oint i A_\mu \dot{x}^\mu ds \right]$$

- Can be defined for an arbitrary curve in spacetime.
- For a pair of antiparallel lines

$$\langle W \rangle \approx \exp \left[ - T V(L, \lambda) \right]$$

- The potential behaves like


$$V(L, \lambda) = \begin{cases} g(\lambda) & \text{screening} \\ \frac{f(\lambda)}{L} & \text{conformal} \\ \alpha' L & \text{confining} \end{cases}$$

- In  $\mathcal{N} = 4$  SYM the most natural Wilson loops includes a coupling to the scalar fields

$$W = \text{Tr} \mathcal{P} \exp \left[ \oint (i A_\mu \dot{x}^\mu + |\dot{x}| n^I \Phi_I) ds \right]$$

$n^I$  do not have to be constant.

- For a smooth loop and continuous  $|n^I| = 1$ , these are finite observables.



## Cusp anomalous dimensions and quark-antiquark potential


- The antiparallel lines suffer also from a subtle linear IR divergence.

## Cusp anomalous dimensions and quark-antiquark potential

- The antiparallel lines suffer also from a subtle linear IR divergence.
- It is simpler to control **logarithmic divergences**.


## Cusp anomalous dimensions and quark-antiquark potential

- The antiparallel lines suffer also from a subtle linear IR divergence.
- It is simpler to control logarithmic divergences.
- Consider Wilson loops with cusps




- All but the black line will suffer from logarithmic divergences.
- Taking  $\phi = i\varphi$  and  $\varphi \rightarrow \infty$  gives the Lorenzian null cusp.

- A compact versions of cusped loops.
- No gauge-invariance subtleties!






- A compact versions of cusped loops.
- No gauge-invariance subtleties!
- Same divergences.
- Completely equivalent, in conformal theories.




- A compact versions of cusped loops.
- No gauge-invariance subtleties!
- Same divergences.
- Completely equivalent, in conformal theories.
- I label the opening angle  $\pi - \phi$ .
- $\phi = 0$  is the circle.
- $\phi \rightarrow \pi$  gives the antiparallel lines.




- A compact versions of cusped loops.
- No gauge-invariance subtleties!
- Same divergences.
- Completely equivalent, in conformal theories.
- I label the opening angle  $\pi - \phi$ .
- $\phi = 0$  is the circle.
- $\phi \rightarrow \pi$  gives the antiparallel lines.
- Can couple the two arcs to two different scalar fields

$$\Phi_1 \quad \text{and} \quad \Phi_1 \cos \theta + \Phi_2 \sin \theta$$



- A compact versions of cusped loops.
- No gauge-invariance subtleties!
- Same divergences.
- Completely equivalent, in conformal theories.
- I label the opening angle  $\pi - \phi$ .
- $\phi = 0$  is the circle.
- $\phi \rightarrow \pi$  gives the antiparallel lines.
- Can couple the two arcs to two different scalar fields

$$\Phi_1 \quad \text{and} \quad \Phi_1 \cos \theta + \Phi_2 \sin \theta$$




- In a conformal theory, by the usual conformal Ward identity

$$\langle W \rangle \sim \frac{1}{d^{2\Delta}}, \quad d = r \frac{\cos \frac{\phi}{2}}{1 - \sin \frac{\phi}{2}}$$

- $\Delta$  is the coefficient of the log divergence.

- By the inverse exponential map we get the gauge theory on  $\mathbb{S}^3 \times \mathbb{R}$
- These are parallel lines on  $\mathbb{S}^3 \times \mathbb{R}$ .




- By the inverse exponential map we get the gauge theory on  $\mathbb{S}^3 \times \mathbb{R}$
- These are parallel lines on  $\mathbb{S}^3 \times \mathbb{R}$ .
- From this last picture we expect

$$\langle W \rangle \approx \exp \left[ -T V(\phi, \theta, \lambda) \right]$$

- In a conformal theory  $T$  is related to divergence at the cusp by the exponential map

$$T = \log \frac{\Lambda_{\text{IR}}}{\Lambda_{\text{UV}}}$$

- Therefore  $V(\phi, \theta, \lambda)$  is the same as  $\Delta$ , the coefficient of the log divergence.




- By the inverse exponential map we get the gauge theory on  $\mathbb{S}^3 \times \mathbb{R}$
- These are parallel lines on  $\mathbb{S}^3 \times \mathbb{R}$ .
- From this last picture we expect

$$\langle W \rangle \approx \exp \left[ -T V(\phi, \theta, \lambda) \right]$$

- In a conformal theory  $T$  is related to divergence at the cusp by the exponential map

$$T = \log \frac{\Lambda_{\text{IR}}}{\Lambda_{\text{UV}}}$$

- Therefore  $V(\phi, \theta, \lambda)$  is the same as  $\Delta$ , the coefficient of the log divergence.
- This  $V(\phi, \theta, \lambda)$  is the generalization of  $V(L, \lambda)$  — the quark-antiquark potential.
- For a conformal theory it has a pole at  $\phi \rightarrow \pi$  and the residue is  $LV(L, \lambda)$ .
- More generally controls **all** log divergences of **all** Wilson loops.
- Needed for a proper renormalization program of Wilson loop operators (and to derive regularized loop equations).



## Local operator insertions

- There is another source of log divergences in Wilson loops:  
Adjoint valued operators inserted into the Wilson loop.



## Local operator insertions

- There is another source of log divergences in Wilson loops:  
Adjoint valued operators inserted into the Wilson loop.
- For example, one operator in the straight line

$$\begin{aligned} W &= \text{Tr} \mathcal{P} \left[ \mathcal{O}(0) \exp \left( \int (iA_\mu \dot{x}^\mu + \Phi^I n^I |\dot{x}|) ds \right) \right] \\ &= \text{Tr} \left[ \mathcal{P} \exp \left( \int_{-\infty}^0 (iA_\mu \dot{x}^\mu + \Phi^I n^I |\dot{x}|) ds \right) \mathcal{O}(0) \mathcal{P} \exp \left( \int_0^\infty (iA_\mu \dot{x}^\mu + \Phi^I n^I |\dot{x}|) ds \right) \right] \end{aligned}$$

- $\mathcal{O}$  is any adjoint operator, *e.g.*,  $F_{23}$ ,  $Z^L$ , etc.

## Local operator insertions

- There is another source of log divergences in Wilson loops:  
Adjoint valued operators inserted into the Wilson loop.
- For example, one operator in the straight line

$$\begin{aligned} W &= \text{Tr } \mathcal{P} \left[ \mathcal{O}(0) \exp \left( \int (iA_\mu \dot{x}^\mu + \Phi^I n^I |\dot{x}|) ds \right) \right] \\ &= \text{Tr} \left[ \mathcal{P} \exp \left( \int_{-\infty}^0 (iA_\mu \dot{x}^\mu + \Phi^I n^I |\dot{x}|) ds \right) \mathcal{O}(0) \mathcal{P} \exp \left( \int_0^\infty (iA_\mu \dot{x}^\mu + \Phi^I n^I |\dot{x}|) ds \right) \right] \end{aligned}$$

- $\mathcal{O}$  is any adjoint operator, *e.g.*,  $F_{23}$ ,  $Z^L$ , etc.
- In a conformal theory, a Wilson loop with two operator insertions at a distance  $d$  will have a VEV

$$\langle W \rangle \sim \frac{1}{d^{2\Delta}}$$

- $\Delta$  is the coefficient of the log divergences — the conformal dimension of the insertions.

## Generalized quark-antiquark potential in $\mathcal{N} = 4$ SYM

- Crucial point: Calculations of  $V(\phi, \theta, \lambda)$  are no harder than for the antiparallel case!

## Generalized quark-antiquark potential in $\mathcal{N} = 4$ SYM

- Crucial point: Calculations of  $V(\phi, \theta, \lambda)$  are no harder than for the antiparallel case!
- Expanding at weak coupling

$$V(\phi, \theta, \lambda) = \sum_{n=1}^{\infty} \left( \frac{\lambda}{16\pi^2} \right)^n V^{(n)}(\phi, \theta)$$


- And at strong coupling

$$V(\phi, \theta, \lambda) = \frac{\sqrt{\lambda}}{4\pi} \sum_{l=0}^{\infty} \left( \frac{4\pi}{\sqrt{\lambda}} \right)^l V_{AdS}^{(l)}(\phi, \theta)$$

## Perturbative calculation

### 1-loop

- Just the exchange of a gluon and scalar field



- This graph is given by the integral


$$\begin{aligned}\partial_\lambda \langle W \rangle \Big|_{\lambda=0} &= \int_{s < t} ds dt \langle (iA_\mu \dot{x}^\mu(s) + |\dot{x}| \Phi^I n^I(s)) (iA_\mu \dot{x}^\mu(t) + |\dot{x}| \Phi^J n^J(t)) \rangle \\ &= \frac{\lambda}{8\pi^2} \int ds dt \frac{-\dot{x}_\mu(s) \dot{x}^\mu(t) + n^I(s) n^I(t)}{|x(s) - x(t)|^2} \\ &= \frac{\lambda}{8\pi^2} \int ds dt \frac{-\cos \phi + \cos \theta}{s^2 + t^2 + 2st \cos \phi} = -\frac{\lambda}{8\pi^2} \frac{\cos \phi - \cos \theta}{\sin \phi} \phi \log \frac{R}{\epsilon}\end{aligned}$$

- Therefore

$$V^{(1)}(\phi, \theta) = 2 \frac{\cos \phi - \cos \theta}{\sin \phi} \phi$$

## Higher order graphs

- Ladder graphs are relatively easy.
- They dominate a funny double-scaled limit where  $\theta \rightarrow i\infty$  with  $\lambda\theta$  fixed. [Correa, Henn  
Maldacena, Sever]
- They are given by harmonic polylogs apparently to all orders. [Henn, Huber]
- Results at weak and strong coupling match.




## Higher order graphs

- Ladder graphs are relatively easy.
- They dominate a funny double-scaled limit where  $\theta \rightarrow i\infty$  with  $\lambda\theta$  fixed. [Correa, Henn  
Maldacena, Sever]
- They are given by harmonic polylogs apparently to all orders. [Henn, Huber]
- Results at weak and strong coupling match.
- Interacting graphs are a bit more complicated.
- At two loops there are bubble graphs and the single cubic vertex graphs.
- they give

$$V_{\text{int}}^{(2)}(\phi, \theta) = -\frac{2}{3}(\pi^2 - \phi^2)V^{(1)}(\phi, \theta)$$

- Full 3 loop answer was also calculated.


[Correa, Henn  
Maldacena, Sever]



## String calculation

[Maldacena] [Rey, Yee] [Drukker  
Gross, Ooguri]

- Within the  $AdS/CFT$  correspondence Wilson loops are calculated by an infinite open string extending to the boundary of  $AdS$ .






## String calculation

[Maldacena][Rey, Yee][Drukker  
Gross, Ooguri]

- Within the  $AdS/CFT$  correspondence Wilson loops are calculated by an infinite open string extending to the boundary of  $AdS$ .
- At the leading order we should find the minimal surface ending on lines separated by  $\pi - \phi$  on the boundary of  $AdS$  and  $\theta$  on  $S^5$ .
- All the string solutions fit inside  $AdS_3 \times S^1$

$$ds^2 = \sqrt{\lambda} (-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\varphi^2 + d\vartheta^2)$$




## String calculation

[Maldacena][Rey, Yee][Drukker  
Gross, Ooguri]

- Within the  $AdS/CFT$  correspondence Wilson loops are calculated by an infinite open string extending to the boundary of  $AdS$ .
- At the leading order we should find the minimal surface ending on lines separated by  $\pi - \phi$  on the boundary of  $AdS$  and  $\theta$  on  $S^5$ .
- All the string solutions fit inside  $AdS_3 \times S^1$

$$ds^2 = \sqrt{\lambda} (-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\varphi^2 + d\vartheta^2)$$

- The equations of motion and classical action can be solved by elliptic integrals.
- $V_{AdS}^{(0)}$  given by a solution of a transcendental equation



## String calculation


[Maldacena][Rey, Yee][Drukker  
Gross, Ooguri]

- Within the  $AdS/CFT$  correspondence Wilson loops are calculated by an infinite open string extending to the boundary of  $AdS$ .
- At the leading order we should find the minimal surface ending on lines separated by  $\pi - \phi$  on the boundary of  $AdS$  and  $\theta$  on  $S^5$ .
- All the string solutions fit inside  $AdS_3 \times S^1$

$$ds^2 = \sqrt{\lambda} (-\cosh^2 \rho dt^2 + d\rho^2 + \sinh^2 \rho d\varphi^2 + d\vartheta^2)$$

- The equations of motion and classical action can be solved by elliptic integrals.
- $V_{AdS}^{(0)}$  given by a solution of a transcendental equation
- Expand around  $\phi = \theta = 0$  the answer is

$$\begin{aligned} V_{AdS}^{(0)}(\phi, \theta) = & \frac{1}{\pi}(\theta^2 - \phi^2) - \frac{1}{8\pi^3}(\theta^2 - \phi^2)(\theta^2 - 5\phi^2) \\ & + \frac{1}{64\pi^5}(\theta^2 - \phi^2)(\theta^4 - 14\theta^2\phi^2 + 37\phi^4) \\ & - \frac{1}{2048\pi^7}(\theta^2 - \phi^2)(\theta^6 - 27\theta^4\phi^2 + 291\theta^2\phi^4 - 585\phi^6) + O((\phi, \theta)^{10}) \end{aligned}$$



## 1-loop determinant

- Complicated fluctuation problem.
- Can be done analytically (implicitly) for either  $\phi = 0$  or  $\theta = 0$ .
- For  $\theta = 0$  and small  $\phi$  we can expand

$$V_{AdS}^{(1)}(\phi, 0) = \frac{3}{2} \frac{\phi^2}{4\pi^2} + \left( \frac{53}{8} - 3\zeta(3) \right) \frac{\phi^4}{16\pi^4} + \left( \frac{223}{8} - \frac{15}{2}\zeta(3) - \frac{15}{2}\zeta(5) \right) \frac{\phi^6}{64\pi^6} \\ + \left( \frac{14645}{128} - \frac{229}{8}\zeta(3) - \frac{55}{4}\zeta(5) - \frac{315}{16}\zeta(7) \right) \frac{\phi^8}{256\pi^8} + O(\phi^{10})$$

## $\phi \rightarrow \pi$ limit

- $V^{(1)}$ ,  $V^{(2)}$ ,  $V_{AdS}^{(0)}$  and  $V_{AdS}^{(1)}$  all have poles at  $\phi = \pi$
- In perturbation theory

$$V(\phi, \theta) \rightarrow -\frac{\lambda}{8\pi} \frac{1 + \cos \theta}{\pi - \phi} + \frac{\lambda^2}{32\pi^3} \frac{(1 + \cos \theta)^2}{\pi - \phi} \log \frac{e}{2(\pi - \phi)} + O(\lambda^3)$$

## $\phi \rightarrow \pi$ limit

- $V^{(1)}$ ,  $V^{(2)}$ ,  $V_{AdS}^{(0)}$  and  $V_{AdS}^{(1)}$  all have poles at  $\phi = \pi$
- In perturbation theory

$$V(\phi, \theta) \rightarrow -\frac{\lambda}{8\pi} \frac{1 + \cos \theta}{\pi - \phi} + \frac{\lambda^2}{32\pi^3} \frac{(1 + \cos \theta)^2}{\pi - \phi} \log \frac{e}{2(\pi - \phi)} + O(\lambda^3)$$

- In the case of  $\theta = 0$  we get essentially the same as the antiparallel lines with  $L \rightarrow \pi - \phi$

$$V(L, \lambda) = \begin{cases} -\frac{\lambda}{4\pi L} + \frac{\lambda^2}{8\pi^2 L} \ln \frac{T}{L} + \dots & \lambda \ll 1 \\ \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L} \left( 1 - \frac{1.3359 \dots}{\sqrt{\lambda}} + \dots \right) & \lambda \gg 1 \end{cases}$$

- The strong coupling calculations also agree in the limit.

## Expansions in small angles

- Consider the expansion of  $V(\phi, \theta, \lambda)$  at small  $\phi$  or  $\theta$

$$\frac{1}{2} \frac{\partial^2}{\partial \theta^2} V(\phi, \theta, \lambda) \Big|_{\phi=\theta=0} = -\frac{1}{2} \frac{\partial^2}{\partial \phi^2} V(\phi, \theta, \lambda) \Big|_{\phi=\theta=0} = \begin{cases} \frac{\lambda}{16\pi^2} - \frac{\lambda^2}{384\pi^2} + \dots & \lambda \ll 1 \\ \frac{\sqrt{\lambda}}{4\pi^2} - \frac{3}{8\pi^2} + \dots & \lambda \gg 1 \end{cases}$$

## Expansions in small angles

- Consider the expansion of  $V(\phi, \theta, \lambda)$  at small  $\phi$  or  $\theta$

$$\frac{1}{2} \frac{\partial^2}{\partial \theta^2} V(\phi, \theta, \lambda) \Big|_{\phi=\theta=0} = -\frac{1}{2} \frac{\partial^2}{\partial \phi^2} V(\phi, \theta, \lambda) \Big|_{\phi=\theta=0} = \begin{cases} \frac{\lambda}{16\pi^2} - \frac{\lambda^2}{384\pi^2} + \dots & \lambda \ll 1 \\ \frac{\sqrt{\lambda}}{4\pi^2} - \frac{3}{8\pi^2} + \dots & \lambda \gg 1 \end{cases}$$

- This quantity was named the bremsstrahlung function  $B(\lambda)$  [ Correa, Henn  
Maldacena, Sever ]
- Calculates the radiation of an accelerated quark.
- Is related to small deformations of BPS Wilson loops and can be calculated exactly

$$B = \frac{1}{2\pi^2} \lambda \partial_\lambda \langle W_\circ \rangle$$

$$\langle W_\circ \rangle = \frac{1}{N} L_{N-1}^1 \left( -\frac{\lambda}{4N} \right) e^{\frac{\lambda}{8N}}$$



## Result so far:

Explicit expressions for these families of Wilson loops at weak and strong coupling.

## Wilson loops and integrability

- We want to apply the tools of integrability to the case of Wilson loops:
  - Find a spin-chain model.
  - Find the all loop scattering (and reflection) matrix
  - Try to solve it exactly.
- This will allow to derive the gauge theory perturbative results from world-sheet techniques.

## Wilson loops and integrability


- We want to apply the tools of integrability to the case of Wilson loops:
  - Find a spin–chain model.
  - Find the all loop scattering (and reflection) matrix
  - Try to solve it exactly.
- This will allow to derive the gauge theory perturbative results from world-sheet techniques.
- Main trick will be to start with the Wilson loop with an arbitrary insertion in it, which will simplify the steps above and at the end remove the insertion.
- In the case of the straight line, after removing the insertion, the operator is  $1/2$  BPS, so no anomalous dimension. So need to know how to treat the cusp.

## string picture

- The string dual of a Wilson loop with an insertion is an excited state of the open string describing the Wilson loop.


## string picture

- The string dual of a Wilson loop with an insertion is an excited state of the open string describing the Wilson loop.




## string picture

- The string dual of a Wilson loop with an insertion is an excited state of the open string describing the Wilson loop.



## string picture

- The string dual of a Wilson loop with an insertion is an excited state of the open string describing the Wilson loop.



- Study the spectrum of open string states all satisfying the same boundary conditions.

- An insertion of  $Z^J$  is described by a string ending along the same curve on the boundary but in the bulk of space rotating around the equator of  $S^5$  with momentum  $J$ .
- An excitation traveling along this string will not know that it's an open string and not the usual  $\text{Tr } Z^J$  vacuum.




- An insertion of  $Z^J$  is described by a string ending along the same curve on the boundary but in the bulk of space rotating around the equator of  $S^5$  with momentum  $J$ .
- An excitation traveling along this string will not know that it's an open string and not the usual  $\text{Tr } Z^J$  vacuum.
- Once it gets to the end of the string we should impose boundary conditions.


- An insertion of  $Z^J$  is described by a string ending along the same curve on the boundary but in the bulk of space rotating around the equator of  $S^5$  with momentum  $J$ .
- An excitation traveling along this string will not know that it's an open string and not the usual  $\text{Tr } Z^J$  vacuum.
- Once it gets to the end of the string we should impose boundary conditions.

## Gauge theory picture

We take the cusped Wilson loop with an adjoint valued operator like  $Z^J$  at the cusp.




- It is clear how to see the appearance of the spin-chain by considering the compact operator in the gauge theory




- In this case the classical dimension is 5.

- It is clear how to see the appearance of the spin-chain by considering the compact operator in the gauge theory



- In this case the classical dimension is 5.
- The bulk hamiltonian is like the usual Minahan-Zarembo spin-chain (Beisert S-matrix  $S_{ab}^{cd}(p_1, p_2) \otimes S_{\dot{a}\dot{b}}^{\dot{c}\dot{d}}(p_1, p_2)$  ).
- Boundary interaction has to be studied separately.

- It is clear how to see the appearance of the spin-chain by considering the compact operator in the gauge theory



- In this case the classical dimension is 5.
- The bulk hamiltonian is like the usual Minahan-Zarembo spin-chain (Beisert S-matrix  $\mathbb{S}_{ab}^{cd}(p_1, p_2) \otimes \mathbb{S}_{\dot{a}\dot{b}}^{\dot{c}\dot{d}}(p_1, p_2)$  ).
- Boundary interaction has to be studied separately.
- The two boundaries interact through wrapping effects at  $O(g^{2(J+1)})$ .
- For  $J = 0$  this is at one-loop.

## All loop reflection matrix and a twist

- The one loop bulk hamiltonian is the same as for closed spin-chains
- The boundary reflection matrix was calculated from Feynman graphs only in the  $SU(2)$  sector.

## All loop reflection matrix and a twist

- The one loop bulk hamiltonian is the same as for closed spin-chains
- The boundary reflection matrix was calculated from Feynman graphs only in the  $SU(2)$  sector.
- To do it to all loops we should use the symmetry:

$$\begin{array}{ccc} \mathfrak{psu}(2, 2|4) & \xrightarrow{Z^J \text{ vacuum}} & \mathfrak{psu}(2|2)_L \times \mathfrak{psu}(2|2)_R \\ \text{boundary } \downarrow & & \downarrow \\ \mathfrak{osp}(4^*|4) & \longrightarrow & \mathfrak{psu}(2|2)_D \end{array}$$

- A single boundary breaks the symmetry to a diagonal  $\mathfrak{psu}(2|2)$ .

## All loop reflection matrix and a twist


- The one loop bulk hamiltonian is the same as for closed spin-chains
- The boundary reflection matrix was calculated from Feynman graphs only in the  $SU(2)$  sector.
- To do it to all loops we should use the symmetry:

$$\begin{array}{ccc}
 \mathfrak{psu}(2, 2|4) & \xrightarrow[Z^J \text{ vacuum}]{} & \mathfrak{psu}(2|2)_L \times \mathfrak{psu}(2|2)_R \\
 \text{boundary } \downarrow & & \downarrow \\
 \mathfrak{osp}(4^*|4) & \longrightarrow & \mathfrak{psu}(2|2)_D
 \end{array}$$

- A single boundary breaks the symmetry to a diagonal  $\mathfrak{psu}(2|2)$ .
- By the usual argument, the boundary reflection matrix should have the same matrix structure as the bulk one

$$\mathbb{R}_{a\dot{a}}^{\dot{b}b}(p) = R_0(p) \hat{S}_{a\dot{a}}^{\dot{b}b}(p, -p)$$



- It replaces  $\mathfrak{psu}(2|2)_L \leftrightarrow \mathfrak{psu}(2|2)_R$  labels.







- Need to determine  
 $R_0(p) = \sigma_B(p)/\sigma(p, -p)$ .
- Like the crossing relation in the bulk, there is a boundary “crossing-unitarity equation”

$$\mathbb{R}(p) = \mathbb{S}(p, -p)\mathbb{R}^c(\bar{p})$$



- Need to determine  $R_0(p) = \sigma_B(p)/\sigma(p, -p)$ .
- Like the crossing relation in the bulk, there is a boundary “crossing-unitarity equation”

$$\mathbb{R}(p) = \mathbb{S}(p, -p)\mathbb{R}^c(\bar{p})$$



- This translates to the conditions on  $\sigma_B$



$$\sigma_B(p)\sigma_B(\bar{p}) = \frac{x^- + 1/x^-}{x^+ + 1/x^+}, \quad \sigma_B(p)\sigma_B(\bar{p}) = 1.$$

where the Joukowski variables are a solution of

$$e^{ip} = \frac{x^+}{x^-}, \quad x^+ + \frac{1}{x^+} - x^- - \frac{1}{x^-} = \frac{1}{g}.$$

- Need to determine  $R_0(p) = \sigma_B(p)/\sigma(p, -p)$ .
- Like the crossing relation in the bulk, there is a boundary “crossing-unitarity equation”

$$\mathbb{R}(p) = \mathbb{S}(p, -p)\mathbb{R}^c(\bar{p})$$



- This translates to the conditions on  $\sigma_B$

$$\sigma_B(p)\sigma_B(\bar{p}) = \frac{x^- + 1/x^-}{x^+ + 1/x^+}, \quad \sigma_B(p)\sigma_B(\bar{p}) = 1.$$

where the Joukowski variables are a solution of

$$e^{ip} = \frac{x^+}{x^-}, \quad x^+ + \frac{1}{x^+} - x^- - \frac{1}{x^-} = \frac{1}{g}.$$

- The solution which matches the all consistency requirements is

$$\sigma_B(z) = \frac{1 + 1/(x^-)^2}{1 + 1/(x^+)^2} e^{-i\chi_B(x^+) + i\chi_B(x^-)}$$

where

$$\chi_B(x) = -i \oint \frac{dz}{2\pi i} \frac{1}{x-z} \log \frac{\sinh 2\pi g(z + 1/z)}{2\pi g(z + 1/z)}.$$

- So far only right boundary. What about the left?


- So far only right boundary. What about the left?
- The left boundary is essentially the same.
- The choice of diagonal subgroup  $\mathfrak{psu}(2|2)_L \times \mathfrak{psu}(2|2)_R \rightarrow \mathfrak{psu}(2|2)_{D'}$  may be different.
- Conjugate the reflection matrix by a twist matrix  $\mathbb{G}$  acting on the  $\mathfrak{psu}(2|2)_L$  labels

$$\mathbb{G} = \text{diag}(e^{i\theta/2}, e^{-i\theta/2}, e^{i\phi/2}, e^{-i\phi/2})$$

- So far only right boundary. What about the left?
- The left boundary is essentially the same.
- The choice of diagonal subgroup  $\mathfrak{psu}(2|2)_L \times \mathfrak{psu}(2|2)_R \rightarrow \mathfrak{psu}(2|2)_{D'}$  may be different.
- Conjugate the reflection matrix by a twist matrix  $\mathbb{G}$  acting on the  $\mathfrak{psu}(2|2)_L$  labels

$$\mathbb{G} = \text{diag}(e^{i\theta/2}, e^{-i\theta/2}, e^{i\phi/2}, e^{-i\phi/2})$$


- This is all the information needed to understand the spectrum of asymptotically large insertions into the Wilson loop.



- So far only right boundary. What about the left?
- The left boundary is essentially the same.
- The choice of diagonal subgroup  $\mathfrak{psu}(2|2)_L \times \mathfrak{psu}(2|2)_R \rightarrow \mathfrak{psu}(2|2)_{D'}$  may be different.
- Conjugate the reflection matrix by a twist matrix  $\mathbb{G}$  acting on the  $\mathfrak{psu}(2|2)_L$  labels

$$\mathbb{G} = \text{diag}(e^{i\theta/2}, e^{-i\theta/2}, e^{i\phi/2}, e^{-i\phi/2})$$

- This is all the information needed to understand the spectrum of asymptotically large insertions into the Wilson loop.




- But not the case  $J = 0 \dots$

## Wrapping effects and the quark-antiquark potential


- One can derive a set of boundary thermodynamic Bethe ansatz equations for this open spin-chain.
- This can be simplified in the small angle limit, where the full answer was reproduced.  
[Correa, Maldacena, Sever] [Gromov Sever]
- They are the same as the usual TBA equations with several small modifications:
  - The  $Y$  functions are related by reflection  $Y_{a,s}(-u) = Y_{a,-s}(u)$
  - There are chemical potentials dependent on  $\phi$  and  $\theta$ .
  - There is a complicated driving term for the massive  $Y_{a,0}$  nodes (aka  $Y_Q$ ).
- The  $Y$ -system equations are unmodified.
  - Analytic properties of the functions are different (determined by the asymptotic solution).




- To reproduce the one loop answer it is enough to consider Lüscher-like corrections.
- This requires to calculate the eigenvalues of the transfer matrix



- To reproduce the one loop answer it is enough to consider Lüscher-like corrections.
- This requires to calculate the eigenvalues of the transfer matrix



- by repeated use of the Yang-Baxter equation this simplifies to



- That is just the product of two **twisted  $\mathfrak{psu}(2|2)$**  transfer matrices.

- On the  $Z^J$  vacuum this is

$$\begin{aligned} T_Q^{\phi, \theta}(p) &= \text{sTr} \left[ \mathbb{R}^{(R)}(p) \mathbb{R}^{(L)c}(\bar{p}) \right] = \text{sTr} \left[ \mathbb{R}^{(R)}(p) \mathbb{G} \mathbb{R}^{(R)c}(-\bar{p}) \mathbb{G} \right] \\ &= \sigma_B(p) \sigma_B(-\bar{p}) \left( \frac{x^-}{x^+} \right)^2 (\text{sTr} \mathbb{G})^2 \end{aligned}$$

- On the  $Z^J$  vacuum this is

$$\begin{aligned} T_Q^{\phi,\theta}(p) &= \text{sTr} \left[ \mathbb{R}^{(R)}(p) \mathbb{R}^{(L)c}(\bar{p}) \right] = \text{sTr} \left[ \mathbb{R}^{(R)}(p) \mathbb{G} \mathbb{R}^{(R)c}(-\bar{p}) \mathbb{G} \right] \\ &= \sigma_B(p) \sigma_B(-\bar{p}) \left( \frac{x^-}{x^+} \right)^2 (\text{sTr} \mathbb{G})^2 \end{aligned}$$

- Simple group theory gives

$$(\text{sTr}_Q \mathbb{G})^2 = 4(\cos \phi - \cos \theta)^2 \frac{\sin^2 Q\phi}{\sin^2 \phi}$$

And the Lüscher-Bajnok-Janik formula is

$$\delta E \approx -\frac{1}{2\pi} \sum_{Q=1}^{\infty} \int_0^{\infty} d\tilde{p} \log \left( 1 + T_Q^{(\phi,\theta)}(\tilde{p}) e^{-2J\tilde{E}_Q} \right)$$

- On the  $Z^J$  vacuum this is

$$\begin{aligned} T_Q^{\phi, \theta}(p) &= \text{sTr} \left[ \mathbb{R}^{(R)}(p) \mathbb{R}^{(L)c}(\bar{p}) \right] = \text{sTr} \left[ \mathbb{R}^{(R)}(p) \mathbb{G} \mathbb{R}^{(R)c}(-\bar{p}) \mathbb{G} \right] \\ &= \sigma_B(p) \sigma_B(-\bar{p}) \left( \frac{x^-}{x^+} \right)^2 (\text{sTr} \mathbb{G})^2 \end{aligned}$$

- Simple group theory gives

$$(\text{sTr}_Q \mathbb{G})^2 = 4(\cos \phi - \cos \theta)^2 \frac{\sin^2 Q\phi}{\sin^2 \phi}$$

And the Lüscher-Bajnok-Janik formula is

$$\delta E \approx -\frac{1}{2\pi} \sum_{Q=1}^{\infty} \int_0^{\infty} d\tilde{p} \log \left( 1 + T_Q^{(\phi, \theta)}(\tilde{p}) e^{-2J\tilde{E}_Q} \right)$$

- Normally for small  $g$  (or large  $J$ ) can expand the logarithm

$$\delta E \approx \frac{1}{2\pi} \sum_{Q=1}^{\infty} \int_0^{\infty} d\tilde{p} T_Q^{(\phi, \theta)}(\tilde{p}) e^{-2J\tilde{E}_Q}$$

For  $J = 0$  the answer will be proportional to  $\frac{g^4(\cos \phi - \cos \theta)^2}{\sin^2 \phi} \dots$

- Crucial fact is that the dressing factor has a double pole at  $\tilde{p} = 0$

$$\begin{aligned}\sigma_B(\tilde{p})\sigma_B(-\tilde{p}) &= e^{2i(\chi_B(x^+) + \chi_B(x^-))} \frac{(2\pi g)^2(x^+ + 1/x^+)(x^- + 1/x^-)}{\sinh(2\pi g(x^+ + 1/x^+)) \sinh(2\pi g(x^- + 1/x^-))} \\ &= e^{2i(\chi_B(x^+) + \chi_B(x^-))} \frac{(2\pi)^2(u^2 + Q^2/4)}{\sinh^2(2\pi u)} \sim \frac{Q^2}{\tilde{p}^2}\end{aligned}$$

- Then using

$$\int_0^\infty d\tilde{p} \log\left(1 + \frac{c}{\tilde{p}^2}\right) = \pi\sqrt{c},$$

- Crucial fact is that the dressing factor has a double pole at  $\tilde{p} = 0$

$$\begin{aligned}\sigma_B(\tilde{p})\sigma_B(-\tilde{p}) &= e^{2i(\chi_B(x^+) + \chi_B(x^-))} \frac{(2\pi g)^2(x^+ + 1/x^+)(x^- + 1/x^-)}{\sinh(2\pi g(x^+ + 1/x^+)) \sinh(2\pi g(x^- + 1/x^-))} \\ &= e^{2i(\chi_B(x^+) + \chi_B(x^-))} \frac{(2\pi)^2(u^2 + Q^2/4)}{\sinh^2(2\pi u)} \sim \frac{Q^2}{\tilde{p}^2}\end{aligned}$$

- Then using

$$\int_0^\infty d\tilde{p} \log\left(1 + \frac{c}{\tilde{p}^2}\right) = \pi\sqrt{c},$$

- The residue is

$$\sqrt{T_Q^{\text{res}} e^{-2J\tilde{E}_Q}} = 2 \frac{\cos\phi - \cos\theta}{\sin\phi} \sin Q\phi (-1)^Q \left[ \frac{(4g^2)^{J+1}}{Q^{2J+1}} - 2(J+2) \frac{(4g^2)^{J+2}}{Q^{2J+3}} + \dots \right]$$

- so

$$\begin{aligned}\delta E &\approx -(4g^2)^{J+1} \frac{\cos\phi - \cos\theta}{\sin\phi} \sum_{Q=1}^{\infty} \frac{(-1)^Q \sin Q\phi}{Q^{2J+1}} \\ &= -\frac{(4g^2)^{J+1}}{2i} \frac{\cos\phi - \cos\theta}{\sin\phi} \left( \text{Li}_{2J+1}(-e^{i\phi}) - \text{Li}_{2J+1}(-e^{-i\phi}) \right)\end{aligned}$$

For  $J = 0$

$$\begin{aligned}\delta E &\approx -\frac{4g^2}{2i} \frac{\cos \phi - \cos \theta}{\sin \phi} (\text{Li}_1(-e^{i\phi}) - \text{Li}_1(-e^{-i\phi})) \\ &= 2g^2 i \frac{\cos \phi - \cos \theta}{\sin \phi} (-\log(1 + e^{i\phi}) + \log(1 + e^{-i\phi})) \\ &= 2g^2 \frac{\cos \phi - \cos \theta}{\sin \phi} \phi + O(g^4)\end{aligned}$$



For  $J = 0$

$$\begin{aligned}\delta E &\approx -\frac{4g^2}{2i} \frac{\cos \phi - \cos \theta}{\sin \phi} (\text{Li}_1(-e^{i\phi}) - \text{Li}_1(-e^{-i\phi})) \\ &= 2g^2 i \frac{\cos \phi - \cos \theta}{\sin \phi} (-\log(1 + e^{i\phi}) + \log(1 + e^{-i\phi})) \\ &= 2g^2 \frac{\cos \phi - \cos \theta}{\sin \phi} \phi + O(g^4)\end{aligned}$$

- This world-sheet calculation is in exact agreement with the one loop perturbative calculation.

For  $J = 0$

$$\begin{aligned}\delta E &\approx -\frac{4g^2}{2i} \frac{\cos \phi - \cos \theta}{\sin \phi} (\text{Li}_1(-e^{i\phi}) - \text{Li}_1(-e^{-i\phi})) \\ &= 2g^2 i \frac{\cos \phi - \cos \theta}{\sin \phi} (-\log(1 + e^{i\phi}) + \log(1 + e^{-i\phi})) \\ &= 2g^2 \frac{\cos \phi - \cos \theta}{\sin \phi} \phi + O(g^4)\end{aligned}$$

- This world-sheet calculation is in exact agreement with the one loop perturbative calculation.
- For Konishi wrapping started at 4 loop order. The cusped Wilson loop is given purely by wrapping from one loop on.
- Is possible to solve iteratively to get higher orders.
- Numerics are hard, but people are working on it.
- Should also be possible to extract the strong coupling answer analytically.

## Summary

When I talked about my paper with Valentina a year ago I would end with the question

Will there be a gauge theory derivation of the strong coupling potential:

$$V(L, \lambda) = \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L}$$

## Summary

When I talked about my paper with Valentina a year ago I would end with the question

Will there be a gauge theory derivation of the strong coupling potential:

$$V(L, \lambda) = \frac{4\pi^2 \sqrt{\lambda}}{\Gamma(\frac{1}{4})^4 L}$$

We are very close to answering **Yes!**

The end